

# EM Scattering

## Homework assignment 2

---

### **Problem 1:**

Consider a dielectric layer in free space. The layer has a relative dielectric constant of  $\epsilon_d = 2.25$ , is infinite, and is parallel to the x-y plane. A plane wave coming from below the layer hits it at an angle equal to  $\theta_i$ . The polarization of the incident wave is either TE<sup>z</sup> where the electric field has no normal component (e.g., in the y-direction) or TM<sup>z</sup> where the magnetic field has no normal component. The frequency of the wave is  $f = 100\text{GHz}$ .

1. Plot the magnitude of the reflection coefficient as function of  $\theta_i$  in both TE and TM cases when the layer is 1 mm-thick
2. Plot the magnitude of the reflection coefficient as function of  $\theta_i$  in both TE and TM cases when the layer is 5 mm-thick

### **Problem 2:**

A plane radio wave is incident on the ionosphere from the air. The relative dielectric constant of the ionosphere is  $\epsilon_d = 1 - (f_p / f)^2$  where  $f$  is the frequency of the wave and  $f_p = 5\text{MHz}$  is the plasma frequency of the ionosphere. The ionosphere is assumed to be infinitely thick.

1. Compute the reflection coefficient for both TE and TM polarizations when the frequency of the wave is  $f = 3\text{MHz}$ .
2. Repeat this calculation at the frequency of  $f = 6\text{MHz}$ . Find the Brewster angle and the critical angle for total reflection.

### **Problem 3:**

Consider the scalar wave equation  $(\nabla^2 + k^2)\psi = 0$  in cylindrical coordinates, and a particular solution given by

$$\psi(\rho, \phi) = \exp(-jm\phi) J_m(k\rho)$$

where  $k$  is the wave number of the medium.

1. Find the corresponding  $\mathbf{M}$  and  $\mathbf{N}$  vectors
2. Sketch the real and imaginary parts of their components along the radial direction (as function of  $\rho$ ) for  $m=1$  and  $\phi = 0$
3. If these vectors are to represent an electric field how does the polarization change as function of radial distance  $\rho$ ? (Again take  $m=1$  and  $\phi = 0$ )

**Problem 4:**


We would like to solve the 2D, **scalar** scattering problem of a scalar wave incident upon an infinitely long circular cylinder whose radius is given by  $a$  and whose axis coincides with the  $z$ -axis. The incident wave is

$$\Psi_i(x, y) = A \exp(-jk_{i,x}x - jk_{i,y}y)$$

with  $k_{i,x}^2 + k_{i,y}^2 = k^2$  where  $k$  is the wave number in the background medium. Like the incident wave, the scattered wave  $\Psi_s(x, y)$  satisfies the 2D wave equation

$$\frac{\partial^2 \Psi_s}{\partial x^2} + \frac{\partial^2 \Psi_s}{\partial y^2} + k^2 \Psi_s = 0$$

1. Using cylindrical coordinates, find the general solution of this equation in terms of a functional series. Take into account the behavior of the scattered wave at infinity.
2. The boundary condition on the surface of the cylinder is  $\Psi = 0$  in which  $\Psi = \Psi_i + \Psi_s$  is the total field. Using this boundary condition, solve the scattering problem by determining the coefficients of the series found above.



This document was created with Win2PDF available at <http://www.win2pdf.com>.  
The unregistered version of Win2PDF is for evaluation or non-commercial use only.  
This page will not be added after purchasing Win2PDF.